

Rituxan Order Form

Pre-filled order form for Rituxan (rituximab) — B-cell depleting therapy for autoimmune disease

REFERRING PROVIDER

PROVIDER NAME	NPI	DEA (IF CONTROLLED)	SPECIALTY
---------------	-----	---------------------	-----------

PRACTICE / CLINIC NAME	OFFICE PHONE	OFFICE FAX
------------------------	--------------	------------

OFFICE CONTACT NAME	CONTACT PHONE	CONTACT EMAIL
---------------------	---------------	---------------

PATIENT INFORMATION

PATIENT LAST NAME	FIRST NAME	MIDDLE INITIAL
-------------------	------------	----------------

DATE OF BIRTH	SEX	PATIENT PHONE	BEST TIME TO CONTACT
---------------	-----	---------------	----------------------

ADDRESS	CITY, STATE, ZIP
---------	------------------

INSURANCE

PRIMARY INSURANCE	MEMBER ID	GROUP #
-------------------	-----------	---------

SECONDARY INSURANCE (IF ANY)	MEMBER ID	GROUP #
------------------------------	-----------	---------

DIAGNOSIS & CLINICAL JUSTIFICATION

PRIMARY DIAGNOSIS	ICD-10 CODE
-------------------	-------------

SECONDARY DIAGNOSIS (IF APPLICABLE)	ICD-10 CODE
-------------------------------------	-------------

Common indications for this medication: *Rheumatoid arthritis, Granulomatosis with polyangiitis (GPA), Microscopic polyangiitis (MPA), Pemphigus vulgaris*

BRIEF CLINICAL SUMMARY / JUSTIFICATION FOR THERAPY

PRIOR THERAPIES TRIED AND OUTCOME
RITUXAN REFERENCE

Generic name: rituximab
Specialty: Rheumatology & Hematology
Typical infusion time: 4–6 hours (first); 2–3 hours (subsequent)
Typical schedule: Two infusions 2 weeks apart, repeated every 6 months
Biosimilars / brands: Truxima, Ruxience, Riabni

MEDICATION ORDER

MEDICATION (DEFAULT: RITUXAN)		GENERIC / ALT (RITUXIMAB)	STRENGTH / FORMULATION
DOSE		FREQUENCY (TYPICAL: TWO INFUSIONS 2 WEEKS APART, REPEATED EVERY 6 MONTHS)	DURATION (TYPICAL: 4&NDASH;6 HOURS (FIRST); 2&NDASH;3 HOURS (SUBSEQUENT))
ROUTE	PRE-MEDICATIONS (IF ANY)		SPECIAL INSTRUCTIONS
ANTICIPATED START DATE	# OF DOSES AUTHORIZED	SUBSTITUTION ALLOWED (Y/N)	BIOSIMILAR ACCEPTABLE (Y/N)

REQUIRED LABS & MONITORING

Check all labs the prescriber requires Ascribe to verify before each infusion:

- | | | |
|--|--|---|
| ☐ CBC w/ differential | ☐ CMP | ☐ Liver function panel |
| ☐ Hepatitis B/C panel | ☐ TB screen (T-SPOT or QFT) | ☐ Pregnancy test (urine) |
| ☐ Iron studies (ferritin, TSAT) | ☐ Vitamin levels (B12, D, folate) | ☐ Other (specify below) |

OTHER LABS / MONITORING REQUIREMENTS
PRESCRIBER SIGNATURE

By signing below I certify that this order is medically necessary, that I am the prescribing provider for this patient, and that I have reviewed the patient's clinical history relevant to this therapy. Ascribe Infusion Therapy is authorized to coordinate scheduling, prior authorization, and benefits verification on the patient's behalf.

PRESCRIBER SIGNATURE
PRINTED NAME
DATE

Form code: ARB-ORDER-RITUXAN · Submit to Ascribe Infusion Therapy by fax to [FAX NUMBER] or via secure portal. Ascribe Infusion will confirm receipt within one business day and begin prior authorization the same day. Questions: [PHONE] · hello@ascribeinfusion.com